

Unusual bacteria causing peritonitis in peritoneal dialysis – A single centre experience

Bactérias raras como causa de peritonite em diálise peritoneal – A experiência de um centro

Ana Farinha, Álvaro Vaz, José Assunção, José Vinhas

Department of Nephrology, Centro Hospitalar de Setúbal, Setúbal, Portugal

Received for publication: 04/04/2012

Accepted in revised form: 18/07/2013

ABSTRACT

Background: Peritonitis is one of the most common complications of peritoneal dialysis. The most frequent agents have been largely described and studied but unusual microorganisms have only been reported in case reports. **Methods:** We describe the cases of unusual bacteria causing peritonitis in our peritoneal dialysis patients between 1998 and 2012. **Results:** In this 15-year period, we have registered 188 peritonitis in 118 patients (0.64 episodes/patient/year). Unusual bacteria were considered all that have caused peritonitis in only one or two instances (12 bacteria in 15 episodes). We have performed antibiogram-based antibiotic therapy in all of them. In two cases catheter removal was necessary and one patient has died in direct relation with the peritonitis. **Conclusions:** Unusual bacterial agents may cause peritonitis in peritoneal dialysis. They are rare and may present with a broad spectrum of seriousness. Publication of experiences may help guiding therapy and increases scientific evidence.

Key-words: Peritoneal dialysis; Peritonitis; Unusual bacteria

RESUMO

Introdução: A peritonite é uma das complicações mais comuns em diálise peritoneal. Os agentes mais frequentes têm sido amplamente descritos e estudados, mas os agentes raros apenas o têm sido em *case reports*. **Métodos:** Descrevemos a casuística de bactérias raras que causaram peritonite nos doentes em diálise peritoneal (DP) da nossa Unidade de 1998 a 2012. **Resultado:** Neste período de 15 anos, registámos 188 peritonites em 118 doentes (0,64 episódios/doente/ano). Foram consideradas bactérias raras todas as que causaram peritonites em um ou dois episódios (12 bactérias em 15 peritonites). A antibioterapia foi baseada no teste de sensibilidade antibiótica de cada microorganismo. Em dois casos foi necessário remover o cateter e noutro caso o paciente morreu em relação directa com a peritonite. **Conclusões:** Algumas

bactérias raras podem causar peritonites em doentes em DP. Estas peritonites são pouco frequentes e podem apresentar uma grande diversidade na sua evolução. A apresentação de experiências aumenta a evidência e pode ajudar a orientar o tratamento destas situações.

Palavras-chave: Bactérias pouco frequentes, Diálise Peritoneal, Peritonite

■ INTRODUCTION

Peritonitis is one of the most frequent complications in peritoneal dialysis and remains one of the most important causes for patient dropout¹. Gram-positives are the most frequent microorganisms responsible for peritonitis². Advances in “connectology” led to a significant decline in the incidence of Gram-positive infections, but new pathogens emerged³. When an unusual agent is present it may be difficult to answer some questions, such as how pathogenic is this agent, how long antibiotics should be prescribed or if the catheter should be removed. Previous case reports may help to guide therapeutic decisions, so we report our single centre experience.

■ RESULTS

Our Unit has followed 118 patients, between 1998 and 2012. Their mean age was 57.7 ± 14.7 years and most were male (53.4%). The end-stage renal disease (ESRD) aetiology was diabetes *mellitus* nephropathy in 30% of cases, chronic glomerulonephritis in 17%, chronic interstitial nephritis in 11%, polycystic disease in 8%, hypertensive nephroangiosclerosis in 7% and in 24% of the cases it was unknown. The mean time in peritoneal dialysis (PD) was 2.5 years. Most of our patients (66%) have been on continuous ambulatory peritoneal dialysis (CAPD) and the remaining were on APD.

We have witnessed a changing pattern in most common agents causing peritonitis in this 15-year period. We registered 188 peritonitis in 118 patients (0.64 episodes/patient/year). Gram-positive were responsible for most peritonitis (50%) but there has been a decrease in *Staphylococcus aureus* incidence from 1.39 episodes/patient/year to 0.1 episodes/patient/year. Gram-negatives were responsible for

21.8% of all peritonitis. *Escherichia coli*, *Pseudomonas aeruginosa* and *Enterobacteriaceae* were the most frequent Gram-negative agents isolated. Mixed infections occurred in 3.2% of the cases and fungi in 2.7%. In 22.3% of the peritonitis no agent has been isolated in effluent cultures.

Our protocol of empirical antibiotherapy has also changed. Between 1998 and 2002, patients diagnosed with peritonitis were treated empirically with intraperitoneal (IP) vancomycin and netilmicin. Since 2002, all patients were treated with intraperitoneal vancomycin and ceftazidime.

Occasionally, we have seen peritonitis by unusual microorganisms. The agents that have caused peritonitis in only one or two instances were considered as unusual (Table 1). Our objective was to describe peritonitis caused by unusual bacteria in our peritoneal dialysis patients and to review the microbiology of these agents. Their clinical situations are summarized in Table 2.

■ ACINETOBACTER LWOFFI

Acinetobacter species are aerobic Gram-negative bacilli that occur in pairs. They are non-fermentative, oxidase-negative and non-motile bacteria (from where its name derives, akinetic)⁴. They can survive on moist and dry surfaces being mainly soil organisms. They can also be found in hospital environments where they may be a source of infection in debilitated patients⁵. *Acinetobacter lwoffii* peritonitis in PD patients has rarely been reported⁶. *Acinetobacter* are innately resistant to many classes of antibiotics, including penicillin, chloramphenicol, and often aminoglycosides. Resistance to fluoroquinolones has also been reported. Carbapenems are recognized as the gold-standard treatment⁷,

Table 1

Unusual bacterial agents causing peritonitis in our Peritoneal Dialysis Unit

Stain	Shape, Characteristics	Genus	Species	Antimicrobial sensitivity pattern	Antimicrobial resistance pattern
Gram +	Coccus	<i>Micrococcus</i>	<i>luteus</i>	Penicillin Ceftriaxone Gentamicin	–
		<i>Micrococcus</i>	<i>spp</i>	Amoxicillin/clavulanic acid Cefuroxime Ciprofloxacin Gentamicin Tetracycline Vancomycin	–
		<i>Kokuria</i>	<i>rosea</i>	Ampicillin Ciprofloxacin Imipenem Rifampin Cotrimoxazol Vancomycin	Ceftazidime
	Bacillus	<i>Corynebacterium</i>	<i>jeikeium</i>	Ampicillin Cefotaxime Netilmicin Tetracycline	Ciprofloxacin Cotrimoxazol
		<i>Brevibacterium</i>	<i>spp</i>	Ampicillin Cefotaxime Netilmicin Vancomycin	–
Gram -	Bacillus, aerobic	<i>Pseudomonas</i>	<i>luteola</i>	Ceftazidime Meropenem	
		<i>Bulkholderia</i>	<i>cepacia</i>	Cotrimoxazol Piperacillin/Tazobactam Meropenem	Cefotaxime Ceftazidime Ceftriaxone Gentamicin Netilmicin
		<i>Stenotrophomonas</i>	<i>maltophilia</i>	Cotrimoxazol Colistin	Cefotaxime Ceftazidime Ceftriaxone Gentamicin Imipenem Netilmicin
		<i>Acinetobacter</i>	<i>lwoffii</i>	Ampicillin Gentamicin Pefloxacin	Cephalothin
		<i>Acinetobacter</i>	<i>lwoffii</i>	Ampicillin Ceftazidime Netilmicin	–
		<i>Rizhobium</i>	<i>radiobacter</i>	Ampicillin Cephalothin Ciprofloxacin Ceftazidime	Amikacin Gentamicin Netilmicin
		<i>Rizhobium</i>	<i>radiobacter</i>	Ampicillin Cephalothin Ciprofloxacin Ceftazidime	–
	Bacillus, facultative anaerobic (Enterobacteriaceae)	<i>Citrobacter</i>	<i>freudii</i>	Cefotaxime Cefuroxime-axetil Ciprofloxacin Gentamicin Cotrimoxazol	Amoxicillin/ clavulanic acid Ampicillin Cephalothin Cefpodoxime
		<i>Klebsiela</i>	<i>oxytoca</i>	Cefotaxime Cefoxithin Ciprofloxacin Imipenem Gentamicin	Ampicillin Cephalothin
		<i>Leclercia</i>	<i>adecarboxylata</i>	Ampicillin Ciprofloxacin Gentamicin Cotrimoxazol	–

Table 2

Summary of clinical cases of peritonitis caused by unusual bacteria in our Peritoneal Dialysis Unit

Patient	Agent	Gender	Age (years)	Aetiology CKD	Modality	Time on PD (months)	Number previous peritonitises	Year of infection	Antibiotic	Days of antibiotic	Outcome
1	<i>Acinetobacter lwoffii</i>	male	29	CIN	CAPD	14	5	2000	Ceftazidime + Netilmicin	21	Recovered
2	<i>Acinetobacter lwoffii</i>	male	62	CGN	CAPD	11	0	1999	Netilmicin	21	Recovered
3	<i>Rhizobium radiobacter</i>	male	47	CGN	CAPD	19	1	2001	Ceftazidime + Ciprofloxacin	21	Recovered
4	<i>Rhizobium radiobacter</i>	male	73	DN	CAPD	9	1	2012	Ceftazidime + Ciprofloxacin	21	Recovered
5	<i>Burkholderia cepacia</i>	male	80	Uncertain	CAPD	14	0	2011	Piperacilin Tazobactam + Cotrimoxazol	21	Removed Catheter
6	<i>Citrobacter freundii</i>	male	65	DN	CAPD	27	0	2006	Ceftazidime + Gentamicin	15 (died before completing therapy)	Deceased
7	<i>Stenotrophomonas maltophilia</i>	female	64	CIN	CAPD	1	1	2006	Cotrimoxazol	21	Removed Catheter
8	<i>Leclercia adecarboxylata</i>	female	77	Uncertain	CAPD	51	1	2011	Ceftazidime	15	Recovered
9	<i>Pseudomonas luteola</i>	female	49	CIN	APD	65	2	2011	Ceftazidime	15	Recovered
10	<i>Klebsiella oxytoca</i>	female	74	HTA	APD	22	0	2009	Ceftazidime	21	Recovered
11	<i>Brevibacterium spp</i>	male	53	Uncertain	CAPD	10	0	2009	Vancomycin	15	Recovered
12	<i>Corynebacterium jeikeium</i>	male	60	DN	APD	32	3	2006	Vancomycin	21	Recovered
13	<i>Kocuria rosea</i>	female	82	HTA	CAPD	7	0	2006	Vancomycin	15	Recovered
14	<i>Micrococcus luteus</i>	male	60	DN	APD	29	1	2006	Vancomycin	21	Recovered
15	<i>Micrococcus spp</i>	male	53	Uncertain	CAPD	8	1	2009	Vancomycin	15	Recovered

APD – automatized peritoneal dialysis; CAPD – continuous ambulatory peritoneal dialysis; CGN – chronic glomerulonephritis; CIN – chronic interstitial nephritis; CKD – chronic kidney disease; DN – diabetic nephropathy; HTA – hypertension.

nevertheless, *Acinetobacter lwoffii* (formerly *Mima Polymorpha*) seems to be more susceptible to antibiotics⁴.

In our Unit we have registered the following two cases:

Patient 1. A 62-year-old male on CAPD for 11 months presented for the first time with abdominal pain and cloudy effluent for 24 hours. The white blood cell count (WBCC) of dialysis fluid showed 3500 WBC/mm³, 90% of neutrophils. *A. lwoffii* was isolated. It was resistant to cephalosporins but sensitive to aminoglycosides. He took 21 days of IP netilmicin with a good response.

Patient 2. A 29-year-old male on CAPD for 14 months was admitted to our hospital complaining of abdominal pain and vomiting for 24 hours. A peritoneal dialysis exchange was performed, showing cloudy effluent (1900 WBC/mm³). His evolution to a septic state forced him to stay in-hospital for 6 days. *A. lwoffii* was isolated and it was resistant to first

generation cephalosporins. He was successfully treated with netilmicin and ceftazidime.

■ RHIZOBIUM RADIOBACTER

Rhizobium (formerly *agrobacterium*) species are aerobic Gram-negative motile bacillus. They are mainly soil organisms. Occasionally, they have been associated with human infections especially in immunocompromised patients with indwelling catheters⁸. In PD patients, a few cases of *Rhizobium* peritonitis have been reported⁹⁻¹³. The outcome seems to be favourable when antibiogram-based therapy is used. Prolonged treatment (21 days)^{12,13} has been associated to better outcome obviating the need for catheter removal⁹. Paucity of data does not allow establishing an antibiotic resistance profile.

We have registered two cases of *Rhizobium radiobacter* peritonitis with more than a decade apart.

Patient 1. A 47-year-old male on CAPD presented to the PD Unit with abdominal pain. He denied cloudy effluent, but when an exchange was made, cytological examination showed 3278 WBC/mm³, predominantly neutrophils. *Rhizobium radiobacter* was identified with a resistance pattern to aminoglycosides. He was medicated with ceftazidime IP for 21 days, but 3 days after completing therapy he presented cloudy effluent again. On this occasion, no agent was found, but, assuming the same aetiology, 15 days of IP ciprofloxacin was prescribed according to the previous antibiogram. The same patient presented a third peritonitis but a Gram+ was identified and a new antibiotic course of vancomycin and amikacin was administered for 21 days. Fortunately, he recovered completely and remained on CAPD for one more year peritonitis-free.

Patient 2. A 73-year-old male on CAPD went to the hospital because of abdominal pain and cloudy effluent. When asked, he recalled an exchange with a damaged dialysis bag. The effluent presented 5000 WBC/mm³, mainly neutrophils. IP ceftazidime was prescribed with good initial response. However, because of the previous patient evolution, we decided to add ciprofloxacin per os when *Rhizobium radiobacter* was isolated. He recovered completely. No other peritonitis was registered.

■ BURKHOLDERIA CEPACIA

Burkholderia cepacia is an aerobic Gram-negative bacillus. It is found in human skin as commensal. It may cause pneumonia in immunocompromised patients with underlying lung disease, such as cystic fibrosis or chronic granulomatous disease¹⁴. It has been rarely associated with peritonitis in PD¹⁵ and it is naturally resistant to many common antibiotics, such as aminoglycosides and polymyxin B. Treatment typically requires an antibiotic association that should include some of the following ceftazidime, doxycycline, piperacillin, meropenem, chloramphenicol and trimethoprim/sulfamethoxazole (cotrimoxazole)¹⁶.

Our patient was an 80-year-old male, immunocompromised because of prostatic cancer. He was medicated with Decapeptyl® and Zometa®. He had

history of a *Proteus mirabilis* exit-site infection, 5 months earlier, but without any previous peritonitis. He presented with abdominal pain, vomiting and cloudy effluent. The cell count of peritoneal fluid was 5200 WBC/mm³ with neutrophil predominance (90%). He was empirically medicated following our protocol (vancomycin + gentamicin). Six days later, the peritoneal microbiology was known and the antibiogram showed resistance to aminoglycosides and cephalosporins. The evolution was ominous, so it was decided to keep the patient hospitalized to remove the catheter. A change in antibiotics was done and IV piperacillin-tazobactam and co-trimoxazole per os were initiated. The patient was transferred to haemodialysis.

■ CITROBACTER FREUNDII

Citrobacter freundii is a facultative anaerobic Gram-negative bacillus. It can be found in soil, water, sewage, food and the intestinal tract of animals and humans. It is classified under the *Enterobacteriaceae*'s family¹⁷. As an opportunistic pathogen, *C. freundii* is responsible for a number of significant opportunistic infections. It is known to cause nosocomial infections of the respiratory tract, urinary tract, blood and many other normally sterile sites¹⁸. A few *Citrobacter freundii* peritonitis in PD have been reported^{19,20}. Various patterns of resistance have been described because of extended-spectrum B-lactamase (ESBL)^{19,20}.

Our patient was a 65-year-old diabetic male, with multiple micro and macrovascular complications. He was admitted to our hospital in a septic status. He complained of abdominal pain. The effluent cell count showed 2670 WBC/mm³, with 87% polymorphonuclear (PMN). Empiric antibiotherapy was initiated. When Gram stain was known (Gram-negative) an abdominal ultrasonography and a CT scan were performed to exclude abdominal complications. *Citrobacter freundii* was isolated and it was resistant to penicillins and first-generation cephalosporins. The antibiotics were changed to ceftazidime and gentamicin following the antibiogram, but the patient maintained cloudy effluent and abdominal pain. Therefore, IV piperacillin-tazobactam was initiated. The patient died before the scheduled catheter removal.

■ *STENOTROPHOMONAS MALTOPHILIA*

Stenotrophomonas maltophilia is an aerobic, non-fermentative Gram-negative bacterium. Initially classified as *Pseudomonas maltophilia*, *S. maltophilia* was also grouped in the genus *Xanthomonas* and, finally, in the genus *Stenotrophomonas*, in 1993. *Stenotrophomonas maltophilia* is catalase and oxidase-negative (distinguishing it from other members)²¹. It is ubiquitous in aqueous environments, soil and plants²². It is not a virulent organism but may be associated to infected prosthesis that may be necessary to remove, like PD catheters. Peritonitis in PD patients with this agent have been reported with and without catheter infection²³⁻²⁶. *Stenotrophomonas maltophilia* is naturally resistant to many broad-spectrum antibiotics (including carbapenems), due to the production of β -lactamases¹⁶. Sensitivity testing requires non-standard culture techniques with incubation at 30 °C²⁷. Testing at the wrong temperature results in isolates being incorrectly reported as being susceptible when they are, in fact, resistant. Many strains of *S. maltophilia* are sensitive to cotrimoxazole and ticarcillin, although resistance has been increasing²⁸. It is not usually susceptible to piperacillin, and susceptibility to ceftazidime is variable²⁹.

We report a 64-year old woman, who presented peritonitis to *Pseudomonas aeruginosa* 10 days after initiating CAPD. After antibiogram was known, she was medicated with ceftazidime and gentamicin but, 10 days later, she complained once more of abdominal pain. The effluent cell count showed 5000 WBC/mm³ 90% PMN, so it was decided to remove the catheter immediately. *Stenotrophomonas maltophilia* was isolated in the peritoneal effluent and in the removed catheter. By that time, antimicrobial sensitivity pattern was known and the patient was treated for 21 days with cotrimoxazol. She was switched to haemodialysis.

■ *LECLERCIA ADECARBOXYLATA*

Leclercia adecarboxylata is a facultative anaerobic Gram-negative bacillus. It is classified under the *Enterobacteriaceae*'s family and formerly known as *Escherichia adecarboxylata*. It is the unique of its genus. Its pathogeny is uncertain and it rarely causes human infection³⁰. Only two cases have been report-

ed in the literature, as far as we know³¹. The antibiotic of the bacillus is unknown.

A 77-year old woman was admitted to our PD Unit for abdominal pain and cloudy dialysate. The WBCC on dialysate showed 6000/mm³, 80% PMN. Empiric antibiotherapy was initiated with vancomycin and ceftazidime IP. When *L. adecarboxylata* was isolated, ceftazidime IP was maintained for 15 days and the patient presented an excellent recovery.

■ *PSEUDOMONAS LUTEOLA*

Pseudomonas luteola is an aerobic, non-fermentative Gram-negative bacillus. In 1987, it was moved to the genus *Chryseomonas luteola*, but returned again to *Pseudomonas* genus in 1997³². It is a soil and water saprophyte but it has been rarely associated with cellulitis and bacteraemia³³. *Pseudomonas luteola* peritonitis in PD has been described previously in one case³⁴. Resistance profiles have been described to amoxicillin, first generation cephalosporin and co-trimoxazole³⁵.

We present a 49-year old woman, with CKD due to staghorn calculi. She had two previous peritonitis to *Pseudomonas putida*, the first two years before and another, one year before. These peritonitis were respectively treated with ceftazidime for 21 days and with ciprofloxacin plus gentamicin for 21 days. She presented a third time with fever (37.8 °C), abdominal pain and cloudy peritoneal dialysate (29.100 WBC/mm³, 90% PMN). On this occasion, *Pseudomonas luteola* was identified and the evolution with IP ceftazidime was uneventful.

■ *KLEBSIELA OXYTOCA*

Klebsiella oxytoca is a facultative anaerobic Gram-negative bacillus from the *Enterobacteriaceae*'s family. It is recognized as a rare human pathogen. To our knowledge, there is only one case of peritonitis in PD caused by *Klebsiella oxytoca* and it was a mixed infection with *Pseudomonas aeruginosa*³⁶. No resistances have been reported.

A 74-year old woman was admitted to our PD Unit for nausea, vomiting and abdominal pain for

48 hours. The dialysate was cloudy and 17500 WBC/mm³, 90% PMN were identified. She initiated vancomycin and ceftazidime. When she was re-evaluated 48 hours later, she was completely asymptomatic. *Klebsiella oxytoca* was isolated, vancomycin was stopped and the peritonitis cured.

■ BREVIBACTERIUM SPECIES

Brevibacterium is a Gram-positive bacillus. Some species are found in cheese. *Brevibacterium casei* and *Brevibacterium epidermidis* are found in human skin and are responsible for odour. Sometimes it is responsible for infections, particularly associated with prosthesis³⁷. A few cases of peritonitis have been described in PD patients³⁸. Resistance to β -lactamics, ciprofloxacin, clindamycin and erythromycin were described and it is sensitive to gentamicin, rifampicin and tetracycline.

A 53-year-old male on CAPD was admitted because of cloudy dialysate and abdominal pain. The WBCC showed 585 WBC/mm³, 85% PMN. A *Brevibacterium* spp was identified and it was sensitive to all tested antibiotics. He was treated with vancomycin with an excellent evolution: in 24 hours he was asymptomatic.

■ MICROCOCCUS SPECIES

Micrococcus is an obligate aerobe Gram-positive coccus that typically appears in tetrads. *Micrococcus* occurs in a wide range of environments, including water, dust, and soil. *Micrococcus* is generally thought to be a saprophytic or commensal organism, although it can be an opportunistic pathogen, particularly in immunocompromised patients, such as those with HIV³⁹. It can be difficult to identify *Micrococcus* as the cause of an infection, since the organism is normally present in skin microflora, and the genus is seldom linked to disease, but recurrent bacteremia, septic shock, septic arthritis, endocarditis, meningitis, and cavitating pneumonia have been reported. A few cases of peritonitis have also been described⁴⁰. *Micrococcus luteus* colonizes the mammalian skin. Moreover, the bacterium colonizes the human mouth, mucosae, oropharynx and

upper respiratory tract. To exclude *Staphylococcus aureus*, a bacitracin susceptibility test can be performed⁴¹.

We had at our Unit two cases of *Micrococcus* peritonitis:

Patient 1. A 60-year-old diabetic male with advanced retinopathy that seriously compromised his vision, had been on APD for 2 years. He was admitted to our Unit due to accidental disconnection of the cyclor lines during the treatment. An exchange with dwell time of 2 hours was done to obtain a dialysate sample. The WBCC showed 880 WBC/mm³, 80% PMN and, therefore, empiric IP antibiotics were administered. A *M. luteus* was isolated and the patient was kept on vancomycin therapy for 21 days on APD with excellent evolution.

Patient 2. The same patient that presented a *Brevibacterium* peritonitis, was readmitted three months latter for cloudy effluent. He denied abdominal pain, as he denied any problem with the PD technique. The WBCC revealed 1900/mm³ with 90% PMN. A *Micrococcus* was isolated. The species was not identified. Vancomycin was maintained for 15 days and the patient remained free from peritonitis for the following 3 years.

■ CORYNEBACTERIUM JEIKEIUM

Corynebacterium jeikeium is an aerobic Gram-positive bacillus. It is pathogenic, typically causing an opportunistic infection, seen most frequently in bone marrow transplant patients but a few cases in PD have been reported⁴²⁻⁴³. It is usually susceptible to vancomycin and tetracycline. Resistance to macrolide antibiotics is common⁴⁴.

Our patient that previously developed *M. luteus* peritonitis was admitted two months later with abdominal pain. The WBCC obtained from dialysate showed 3900/mm³, 80% PMN. By that time no risk factor has been identified. *Corynebacterium jeikeium* was isolated and the patient was successfully treated with vancomycin for 21 days. Like in other unusual Gram-positive peritonitis, the evolution was excellent. The patient was kept in APD at his request.

■ KOCURIA ROSEA

Kocuria rosea is an aerobic, gram-positive coccus. It is found in tetrads, irregular clusters or cubical packets of eight. It was previously classified into the genus *Micrococcus*. This organism is widespread in nature and is frequently found as normal skin flora in humans and other mammals. In this genus, the majority of strains are non-pathogenic, but *Kocuria rosea* may be opportunistic. It has been reported to cause catheter-related bacteraemia⁴¹ and peritonitis⁴⁵⁻⁴⁶. Apart from nitrofurantoin, no other resistances have been reported⁴⁷.

An 82-year-old woman was admitted to our hospital because of severe abdominal pain and cloudy peritoneal dialysate for 5 hours. The WBCC in effluent showed $> 16000/\text{mm}^3$, $> 95\%$ PMN. A *K. rosea* was isolated a few days later. Contrasting with the severity of presentation, after two dwells with empiric antibiotic the patient was asymptomatic. She took 15 days of IP vancomycin with complete recovery.

■ DISCUSSION

In 15 years we have registered 15 peritonitis due to unusual bacterial agents in 12 patients (0.05 episodes/patient/year). Twelve patients recovered completely without need for catheter removal, but in two cases (*Burkholderia cepacia* and *Stenotrophomonas maltophilia*) there was a need to drop-out of haemodialysis, and one patient died in direct relation to his peritonitis (*Citrobacter freundii*).

The International Society of Peritoneal Dialysis (ISPD) infection guidelines advise to deliver antibiotic therapy according to the resistance pattern (possibly with 2 drugs) for 14-21 days, if a clinical response is seen, but the catheter must be removed if no clinical improvement is achieved within 5 days⁴⁸.

In our protocol, we do 2 weeks of antibiotic for most common bacterial peritonitis and 3 weeks for *Staphylococcus aureus*, *Pseudomonas species*, *Xanthomonas species* and other Gram-negatives. In general, Gram-negative peritonitis precludes longer treatments, because they usually have worse prognosis. We applied that principle to our Gram-negative peritonitis presented here, so we do not know if shorter regimens could be effective.

Until enough data about unusual bacteria peritonitis exists, it is wise to individualize treatment and monitor the response to treatment. Catheter removal should be proposed if the organism is multi-resistant or if the clinical situation deteriorates.

In our opinion, a national PD registry would help us defining what is uncommon, how to treat unusual peritonitis depending on patterns of resistance and what to expect from each of those rare situations. The therapeutic strategy for rare peritonitis agents in PD can be improved with new published experiences.

Conflicts of interest statement: None declared.

References

1. Fried LF, Bernardini J, Johnston JR, Piraino B. Peritonitis influences mortality in peritoneal dialysis patients. *J Am Soc Nephrol* 1996;7:2176-2182
2. Piraino B, Bernardini J, Sorkin M. A five-year study of the microbiologic results of exit site infections and peritonitis in continuous ambulatory peritoneal dialysis. *Am J Kidney Dis* 1987; 10:281-286.
3. Port FK, Held PJ, Nolph KD, Turenne MN, Wolfe RA. Risk of peritonitis and technique failure by CAPD connection technique: a national study. *Kidney Int.* 1992;42(4):967-974.
4. Gerner-Smidt P, Frederiksen W. Acinetobacter in Denmark: I. Taxonomy, antibiotic susceptibility, and pathogenicity of 112 clinical strains. *APMIS* 1993; 101(11):815-825
5. Visca P, Seifert H, Towner KJ. Acinetobacter infection – an emerging threat to human health. *IUBMB Life* 2011;63(12):1048-1054
6. Huddam B, Koçak G, Azak A, Duranay M. *Acinetobacter lwoffii* peritonitis in a patient receiving continuous ambulatory peritoneal dialysis. *Ther Apher Dial* 2013;17(1):117-119.
7. Rahal JJ. Novel antibiotic combinations against infections with almost completely resistant *Pseudomonas aeruginosa* and *Acinetobacter* species. *Clin Infect Dis* 2006; 43 (Suppl 2): S95-99
8. Lai CC, Teng LJ, Hsueh PR, et al. Clinical and microbiological characteristics of Rhizobium radiobacter infections. *Clin Infect Dis* 2004;38(1):149-153
9. Han KH, Han SY. A Case of Rhizobium Radiobacter Peritonitis Cured without Removal of the PD Catheter in a Patient on CAPD. *Korean J Nephrol* 2007;26:634-636
10. Rodby RA, Glick EJ. Agrobacterium radiobacter peritonitis in two patients maintained on chronic peritoneal dialysis. *Am J Kidney Dis* 1991;18(3):402-405
11. Rothe H, Rothenpieler U. Peritonitis due to multiresistant rhizobium radiobacter. *Perit Dial Int* 2007;27(2):214-215
12. Minguela JJ, de-Pablos M, Castellanos T, Ruiz-de-Gauna R. Peritonitis by Rhizobium radiobacter. *Perit Dial Int* 2006;26(1):112
13. Marta R, Dâmaso C, Silva JE, Almeida M. Peritonitis due to Rhizobium radiobacter. *Einstein* 2011; 9:389-390
14. Mahenthalingam E, Urban TA, Goldberg JB. The multifarious, multireplicon *Burkholderia cepacia* complex. *Nat Rev Microbiol* 2005;3(2):144-156.
15. Kang HM, Park JE, Lee HD, et al. A case of Burkholderia cepacia peritonitis in a patient on CAPD. *Korean J Med* 2006;7:219-223

16. McGowan JE Jr. Resistance in nonfermenting gram-negative bacteria: multidrug resistance to the maximum. *Am J Infect Control* 2006;34(5 Suppl 1): S29–37;S64–73
17. Wang JT, Chang SC, Chen YC, Luh KT. Comparison of antimicrobial susceptibility of *Citrobacter freundii* isolates in two different time periods. *J Microbiol Immunol Infect* 2000;33(4):258–262
18. Whalen JG, Mully TW, English JC 3rd. Spontaneous *Citrobacter freundii* infection in an immunocompetent patient. *Arch Dermatol* 2007;143(1): 124–125
19. Gursu M, Aydin Z, Pehlivanoglu F, *et al.* *Citrobacter* peritonitis: two cases and review of the literature. *Perit Dial Int* 2011;31(4):409–411
20. Dervisoglu E, Yumuk Z, Yegenaga I. *Citrobacter freundii* peritonitis and tunnel infection in a patient on continuous ambulatory peritoneal dialysis. *J Med Microbiol* 2008; 57(1):125–127
21. Palleroni N, Bradbury JF. *Stenotrophomonas*, a new bacterial genus for *Xanthomonas maltophilia* (Hugh 1980) Swings *et al.* 1983. *Int J Syst Bacteriol* 1993;43(3): 606–609
22. Denton M, Kerr KG. Microbiological and clinical aspects of infection associated with *Stenotrophomonas maltophilia*. *Clin Microbiol Rev* 1998;11(1):57–80
23. Azak A, Kocak G, Huddam B, Scan G, Duranay M. An unusual cause of continuous ambulatory peritoneal dialysis-associated outpatient peritonitis: *Stenotrophomonas maltophilia*. *Am J Infect Control* 2011;39(7):618
24. Taylor G, McKenzie M, Buchanan-Chell M, Perry D, Chui L, Dasgupta M. Peritonitis due to *Stenotrophomonas maltophilia* in patients undergoing chronic peritoneal dialysis. *Perit Dial Int* 1999;19(3):259–262
25. Tzanetou K, Triantaphyllis G, Tsoutsos D, *et al.* *Stenotrophomonas maltophilia* peritonitis in CAPD patients: susceptibility to antibiotics and treatment outcome: a report of five cases. *Perit Dial Int* 2004;24(4):401–404
26. Baek JE, Jung EY, Kim HJ *et al.* *Stenotrophomonas maltophilia* infection in patients receiving continuous ambulatory peritoneal dialysis. *Korean J Intern Med* 2004;19:104–108
27. Wheat PF, Winstanley TG, Spencer RC. Effect of temperature of antimicrobial susceptibilities of *Pseudomonas maltophilia*. *J Clin Pathol* 1985; 38(9):1055–1058.
28. Pankuch GA, Jacobs MR, Applebaum PC. Susceptibilities of 123 *Xanthomonas maltophilia* strains to ciprofloxacin, PD131628, PD138312, PD140248, ciprofloxacin, and ofloxacin. *Antimicrob Agents Chemother* 1994;38(2): 369–370
29. Pankuch GA, Jacobs MR, Rittenhouse SF, Appelbaum PC. Susceptibilities of 123 strains of *Xanthomonas maltophilia* to eight beta-lactams (including beta-lactam-beta-lactamase inhibitor combinations) and ciprofloxacin tested by five methods. *Antimicrob Agents Chemother* 1994;38(10):2317–2322.
30. Hess B, Burchett A, Huntington MK. *Leclercia adedecarboxylata* in an immunocompetent patient. *J Med Microbiol* 2008;57:896–898
31. Fattal O, Deville JG. *Leclercia adedecarboxylata* peritonitis in a child receiving chronic peritoneal dialysis. *Pediatr Nephrol* 2000;15(3–4):186–187.
32. Casalta JP, Fournier PE, Habib J, Riberi A, Raoult D. Prosthetic valve endocarditis caused by *Pseudomonas luteola*. *BMC Infect Dis* 2005;5:82
33. Arnold FW, Sciortino CV, Riede KA. New Associations With *Pseudomonas luteola* Bacteremia: A Veteran With A History Of Tick Bites And A Trauma Patient With Pneumonia. *The Internet Journal of Infectious Diseases*. 2005;4
34. Amber IJ, Reimer LG. *Pseudomonas* sp. group Ve-2 bacterial peritonitis in a patient on continuous ambulatory peritoneal dialysis. *J Clin Microbiol* 1987;25(4):744–745
35. Chihab W, Alaoui AS, Amar M. *Chryseomonas luteola* identified as the source of serious infections in a Moroccan University Hospital. *J Clin Microbiol* 2004;42(4):1837–1839
36. Yang SB, Yang WC, Chen TW, Lin CC. *Klebsiella oxytoca* and polymicrobial infection in peritoneal dialysis-related peritonitis. *Perit Dial Int* 2004;24(2):196–197
37. Ulrich S, Zbinden R, Pagano M, Fischler M, Speich R. Central venous catheter infection with *Brevibacterium* sp. in an immunocompetent woman: case report and review of the literature. *Infection* 2006;34(2):103–106
38. Choi JS, Kim CS, Park JW, ET AL. Consecutive episodes of peritonitis in a patient undergoing peritoneal dialysis caused by unusual organisms: *Brevibacterium* and *Pantoea agglomerans*. *Kidney Res Clin Pract* 2012; 31:121–123
39. Smith KJ, Neafie R, Yeager J, Skelton HG. *Micrococcus folliculitis* in HIV-1 disease. *Br J Dermatol* 1999;141(3):558–561
40. Kao CC, Chiang CK, Huang JW. *Micrococcus* species-related peritonitis in patients receiving peritoneal dialysis. *Int Urol Nephrol* 2012 Oct 9. [Epub ahead of print]
41. Schiffli H, Lang S. Non-diphtheria corynebacteria and CAPD infections. *Nephrol Dial Transplant* 2009; 24(12):3896–3897
42. Szeto CC, Chow KM, Chung KY, Kwan BC, Leung CB, Li PK. The clinical course of peritoneal dialysis-related peritonitis caused by *Corynebacterium* species. *Nephrol Dial Transplant* 2005;20(12):2793–2796
43. Rosato AE, Lee BS, Nash KA. Inducible macrolide resistance in *Corynebacterium jeikeium*. *Antimicrob Agents Chemother* 2001;45(7):1982–1989
44. Altuntas F, Yildiz O, Eser B, Gundogan K, Sumerkan B, Cetin M. Catheter-related bacteremia due to *Kocuria rosea* in a patient undergoing peripheral blood stem cell transplantation. *BMC Infect Dis* 2004;4(1):62.
45. Dotis J, Printza N, Papachristou F. Peritonitis attributable to *Kocuria rosea* in a pediatric peritoneal dialysis patient. *Perit Dial Int* 2012;32(5):577–578.
46. Kaya KE, Kurtoglu Y, Cesur S, *et al.* Peritonitis due to *Kocuria rosea* in a continuous ambulatory peritoneal dialysis case. *Mikrobiyol Bul* 2009;43(2):335–337
47. Savini V, Catavittello C, Masciarelli G, *et al.* Drug sensitivity and clinical impact of members of the genus *Kocuria*. *J Med Microbiol* 2010;59(Pt12):1395–1402.
48. Li PK, Szeto CC, Piraino B, *et al.* Peritoneal dialysis-related infections recommendations: 2010 update. *Perit Dial Int* 2010;30(4):393–423

Correspondence to:

Dr^a Ana Farinha

Av. D. João II, nº12, 7^a Dto, 2910-548 Setúbal, Portugal

E-mail: alpfarinha@yahoo.com.br